

# SETI@BERKELEY: RADIO, IR AND OPTICAL SEARCHES FOR ALIEN TECHNOLOGY



#### SETI@home

Launched in 1999, SETI@home has engaged over 10 million people in 226 countries in the search for life on other worlds. Currently, participants in the project are generating a collective equivalent of 500 TeraFLOPs/sec. SETI@home II users have performed over  $1.4 \times 10^{22}$  FLOPs to-date. This immense and growing computing power is being used to conduct a commensal sky survey for narrow-band signals utilizing Arecibo's ALFA multi-beam receiver.




NAIC Arecibo Observatory

#### SETI@HOME II Specs 50 MHz

Bandwidth: Spectral **Resolution:** Channels: Sky Coverage: Sensitivity:

0.037-1221 Hz 7 x 10<sup>8</sup> 25% 1.4 x 10<sup>-26</sup> W/m<sup>2</sup>




SETI@Home Screenshot


BWRC



AstroPulse is a companion project to SETI@Home. Rather than looking for a large amount of energy in a narrow frequency range, AstroPulse looks for a large amount of energy in a narrow time slice - a radio pulse. These radio pulses could come from dead stars, massive objects colliding with one another or perhaps a beacon from a very advanced technology.



The plot above shows several radio pulses that have been detected by AstroPulse volunteers. Many of the pulses are clustered within the plane of our own galaxy, which is a good indication that the source of these pulses is not terrestrial. The dashed lines indicate the part of the sky viewable from the Arecibo Observatory. Follow-up observations of these detections are underway.



AST(RON


NASA

ANDREW SIEMION - UC BERKELEY, ASTRON, RADBOUD UNIVERSITY

## AstroPulse

## SERENDIP

SERENDIP is an acronym that stands for the Search for Extraterrestrial Radio Emission from Nearby Developed Intelligent **P**opulations.



**SERENDIP Progress: Past, Present and Future** 

The SERENDIP V.v sky survey uses the 7beam Arecibo L-band Feed Array (ALFA) at Arecibo Observatory to carry out a high sensitivity survey of the sky visible to Arecibo. This survey searches for narrowband signals in an approximately 300 MHz wide band surrounding 1420 MHz. The observations are 10 times more sensitive and 100 times more comprehensive than the most sensitive existing SETI large-area surveys (SERENDIP IV and SETI@home).



ALFA Multi-Beam Receiver

300 MHz

I.5 Hz

2 x 10<sup>9</sup>

I x 10<sup>-25</sup> W/m<sup>2</sup>

Serendip V.v Specs Bandwidth: Spectral Resolution Channels: Sky Coverage: 25% Sensitivity:

The two billion channel SERENDIP V.v spectrometer, deployed at Arecibo in late 2008, analyzes signals from all seven beams simultaneously in dual polarization, effectively observing seven points on the sky simultaneously.

SETI HOME

Radboud University Nijmegen


1.4 x 10<sup>-26</sup> W/m<sup>2</sup>

#### IR Excess

The UC Berkeley Infrared Excess Search attempts to identify excess thermal infrared radiation emitted from highly advanced civilizations by searching the IRAS and 2MASS catalogs and performing follow-up observations on likely candidates with the SETI@home, Serendip V.v, and Optical SETI databases.


The initial dataset consists of 1000 stars with ages determined by Ca II absorption to be > I gigayear to reduce the likelihood of an associated

protoplanetary disk.



Cut-away Diagram of an Idealized Dyson Surface

Infrared excess was determined via the K-[12] indicator. The K band is thought to be quite insensitive to stellar atmosphere fluctuations, and hence provides a stable benchmark for color excess.



Infrared excess for all 540 target stars (upper left) and based on stellar type.

33 IR excess candidates were found and subsequently searched for anomalous radio emission nanosecond-scale optical pulses. No convincing signals were found.


CASPER

## **Optical SETI**

The Optical SETI program at UC Berkeley searches for nanosecond time scale pulses, perhaps transmitted by a powerful pulsed laser operated by a distant civilization. Began in 1997, this program was the first search of its kind.

The pulse search utilizes Berkeley's 30 inch automated telescope at Leuschner Observatory in Lafayette (15 miles east of campus) using a custom built photometer designed specifically for optical Statudth: Spectral Resolution Channels: Sensitivity: PMT 1 PMT 2 Bandwidth:





Schematic Diagram of SETI Photometer

#### **Optical SETI Specs**

Band: Average Flux Limit: Peak Flux Limit: Stars Surveyed: Galaxies Surveyed:

CENTER FOR ASTRONOMY SIGNAL PROCESSING AND ELECTRONICS RESEARCH