TESS discovery of a super-Earth and two sub-Neptunes orbiting the bright, nearby, Sun-like star HD 22946

Luca Cacciapuoti^{1,2}, Laura Inno^{3,4}, Giovanni Covone^{1,4,5}, Veselin B. Kostov⁶, Thomas Barclay^{6,42}, Elisa V. Quintana⁶, Knicole D. Colon⁶, Keivan G. Stassun^{7,8}, Benjamin Hord⁹, Steven Giacalone¹⁰, Stephen R. Kane¹¹, Kelsey Hoffman¹², Jason Rowe¹³, Gavin Wang¹⁴, Kevin I. Collins¹⁵, Karen A. Collins¹⁶, Thiam-Guan Tan^{17, 39}, Francesco Gallo¹, Christian Magliano¹, Riccardo M. Ienco¹, Markus Rabus¹⁸, David R. Ciardi¹⁹, Elise Furlan¹⁹, Steve B. Howell²¹, Crystal L. Gnilka²¹, Nicholas J. Scott²¹, Kathryn V. Lester²¹, Carl Ziegler²², César Briceño²³, Nicholas Law²⁴, Andrew W. Mann²⁴, Christopher J. Burke²⁵, Samuel N. Quinn²⁶, Angelo Ciaramella³, Pasquale De Luca³, Stefano Fiscale³, Alessandra Rotundi³, Livia Marcellino³, Ardelio Galletti³, Ida Bifulco³, Fabrizio Oliva²⁷, Alton

Law²⁴, Andrew W. Mann²⁴, Christopher J. Burke²⁵, Samuel N. Quinn²⁶, Angelo Ciaramella³, Pasquale De Luca³, Stefano Fiscale³, Alessandra Rotund¹, Livia Marcellino³, Ardelio Galletti³, Ida Bifulco³, Fabrizio Oliva²⁷, Alton Spencer²⁸, Lisa Kaltenegger²⁹, Sott McDermott⁹, Zahra Essack^{31,32}, Jon M. Jenkins²¹, Bill Wohler^{12,33}, Joshua N. Winn³⁴, S. Seager^{25, 35, 36}, Roland Vanderspek²⁵, George Zhou³⁷, Avi Shporer³⁸, Diana Dragomir⁴⁰, and William Fong⁴¹
(Affiliations can be found after the references) September 21, 2022
MBTRAT We report the Transiting Exoplanet Survey Satellite (TESS) discovery of a three-planet system around the bright Sun-like star HD 22946 (V≈8.3 mag), also known as TIC 10099000, located 63 parsees away. The system was observed by TESS in sectors 3, 4, 30 and 31 and two planet candidates, labeled TESS Tonsi Singmak and use follow-up burgers of a dwit of dwite system was observed by TESS transi Singmak and use follow-up by TFOP allowed us to constrain the stellar parameters of TO1411, including its radius of 1.15720.025*R*, while planet c and *d* are sub-Neptomian planets, with radii of 2.74 ± 0.14 *R*, and 3.23 ± 0.19 *R*, respectively. By using dynamical simulations, *expensionel*, we indemind the system is dynamically stable and potentially unpacked, with enough space to host at least one more planet berveen *c* and *d*. Finally, given that the star is hright and nearly, ved discuss possibilities for detailed to measurements for Mars-like bodies (e.g., L98-59 b, Kostov et al. 2017). Transiting planet vity, systems soround nearly we discuss possibilities for detailed mass characterization of ther sumospheres with the *d*. *Lost C*, *R*, *e*, 4.75) and sub-Neptunes (1.75 < *R*₈ < 3.5), according to the classification criteria defined by Fulton et al. 2017. Nate et al. 2017; Wint et al. 2017. Wint et al. 2

bright stars offer the opportunity for in-depth observational studies to characterize different planets formed in a common environment. The Transiting Exoplanet Survey Satellite (TESS; Ricker et al. 2015) is a space-borne NASA mission launched in 2018 to survey the sky for transiting exoplanets around these stars. To date, it has contributed to the discovery of more than 200 exoplanets in the Galaxy, with more than 5700 candidates on hold for confirmation¹. These discoveries include planets of differ-

their mass via Doppler spectroscopy and the determination of their atmospheric proprieties through transmission or emission spectroscopy. Besides, multiplanetary systems offer the additional opportunity to perform comparative exoplanetology studies, since we can constrain the formation and evolution of a group of planets that share a common origin, given the fixed proprieties of the host star. In fact, for these systems, we can gather enough information to constrain the orbits and dynamics of the system, and hence to infer the underlying formation and evolution scenarios, see e.g., Ragozzine & Holman (2010); Lissauer et al. (2011); Fabrycky et al. (2014).

¹ See the NASA Exoplanet Archive at https:// exoplanetarchive.ipac.caltech.edu/

In this work, we report the TESS discovery of a multiplanetary system consisting of a Super-Earth and two sub-Neptune-sized planets orbiting the bright (V \approx 8.3 mag), nearby (63 pc), Sun-like star HD 22946, indicated in the TESS Input Catalog (Stassun & Torres 2018) as TIC 100990000 and as TOI 411 in the TESS Objects of Interest list on ExoFOP-TESS².

Sub-Neptunes and super-Earths are of particular interest for planetology studies because, while they represent the highest fraction of planets detected in the Galaxy, they are not observed in the Solar System. The detection and in-depth characterization of such planets orbiting Sun-like stars can help us constrain the formation and evolution pathways along which they form and gain a better understanding of different planetary systems (Bitsch 2019). For example, a depression in the distribution of exoplanets radii between 1.5 and 2 R_{\oplus} (Fulton et al. 2017; Owen & Wu 2017) seems to indicate that these exoplanets accreted an atmosphere at the time of formation that progressively thinned out due to heat-driven processes, such as stellar radiation photoevaporation (Fulton & Petigura 2018) or core-powered mass loss (Ginzburg et al. 2018). Moreover, Loyd et al. (2020) found that doubling the known sample of exoplanets with radii $R < 4R_{\oplus}$ is required to confirm or exclude the role of photo-evaporation in the supposed atmospheric loss of these worlds. Thus, by studying planets in this range, we can shed light on the physical processes that can lead to the loss of a primary atmosphere and the possible accretion of a heavier secondary one.

The paper is organized as follows. In Section 2 we characterize the host star, in Sect. 3 we present the detection of the three transiting candidates from TESS observations and their characterization. In Sect. 4 we analyze the follow-up data to validate their planetary nature. In Sect. 5, we explore the dynamics of the system and the potential for atmospheric characterization of its planets. Finally, in Sect. 6, we summarize our findings.

2. Characterization of the host star

Characterizing the physical proprieties of the host star, such as its mass, M_{\star} , radius, R_{\star} and effective temperature $T_{\rm eff}$, is propaedeutic to constraining the corresponding properties of its planetary system. Therefore, we first focus on determining these quantities for HD 22946, together with its spectral class, surface gravity, log g, metallicity, [Fe/H], and sky-projected rotational velocity, $v \sin i$.

HD 22946 is a bright ($V \approx 8.3 \text{ mag}$) main-sequence star located in the Southern Hemisphere, characterized by a high proper motion and some photometric variability. Basic stellar information is given in 1.

2.1. Stellar variability

In order to understand better the photometric variability of the star, we searched for publicly-available light curves of HD 22946, we only found multi-epoch photometry from the Hipparcos Epoch Photometry catalog (Hip 1997), which includes a hundred measurements over about 3 years, with 3 to 10 measurements per night each month. ³




Fig. 1. Light curve of HD 22946 from the Hipparcos Epoch Photometry data (V- band, top panel), and the corresponding power spectrum (bottom panel). In the top panel, the mean luminosity and standard deviation are also shown: the scatter in the photometric data is compatible with a low-amplitude variability. The peak of the power spectrum, indicated by the orange vertical line, is found at ≈ 0.073 days, and it is likely due to instrumental aliasing.




Fig. 2. Spectral energy distribution of HD 22946. The blue symbols and their vertical error bars represent the photometric measurements available for this source and listed in Table 1. The horizontal bars indicate instead the width of the passband for each data point. The overplotted black line is the best fit model, which allows us to derive the stellar parameters.

The light curve from the Hipparcos data shows a small amplitude variation, with a maximum-minimum difference of about ≈ 0.1 mag, or six times the standard deviation σ of the measurements, as can be seen in Figure 1, where we also show the computed Lomb-Scargle (Lomb 1976; Scargle 1982) periodogram using a Nyquist factor of 10 following recommendation from Richards et al. (2012) and VanderPlas (2018). The peak of the power spectrum is found at around 0.073 days, close to the 0.076 days data gaps of Hipparcos observations (Percy et al. 2002), therefore it is likely due to aliasing. Indeed, the star is classified as static (class "C") in the Hipparcos catalog.

The TESS light curve of HD 22946 also shows mild signs of variability, especially in sectors 30 and 31, as we further discuss in the next section.

² https://exofop.ipac.caltech.edu/tess/

³ Light curves for this star are available in ASAS-SN Sky Patrol database (Shappee et al. 2014; Jayasinghe et al. 2021, and references therein) and in ASAS (Pojmanski 1997), but the stellar magnitude is close to the surveys' saturation limit so we discarded this data. Epoch-photometry from Gaia DR3 is not available for this star.

Table 1. Summary of stellar data used in this work.

Parameters	Value	Source
Name TIC ID	TOI-411 100990000	Guerrero & TESS Science Office (2021) TICv8.1
Alt. name	HD 22946	
	Astrometric P	roperties
α R.A. (hh:mm:ss)	03:39:16.761	Gaia EDR3
δ Dec. (dd:mm:ss)	-42:45:45.185	Gaia EDR3
$\mu_{\alpha} (\text{mas yr}^{-1})$	-51.618 ± 0.015	Gaia EDR3
μ_{δ} (mas yr ⁻¹)	-110.546 ± 0.021	Gaia EDR3
Barycentric RV (km s^{-1})	16.91 ± 0.15	Gaia DR2
Distance (pc)	62.87 ± 0.05	Gaia EDR3
	Photometri	c Data
TESS (mag)	7.757 ± 0.006	TIC v8.1
FUV (mag)	19.45 ± 0.16	GALEX GR6
NUV (mag)	13.014 ± 0.005	GALEX GR6
B_T (mag)	8.903 ± 0.017	Tycho-2
V_T (mag)	8.318 ± 0.012	Tycho-2
G-band (mag)	8.1281 ± 0.0003	Gaia DR2
G_{BP} -band (mag)	8.4206 ± 0.0003	Gaia DR2
G_{RP} -band (mag)	7.7173 ± 0.0003	Gaia DR2
<i>B</i> (mag)	8.79 ± 0.03	TIC v8.1
<i>V</i> (mag)	8.27 ± 0.03	TIC v8.1
J(mag)	7.25 ± 0.04	2MASS
H (mag)	7.04 ± 0.04	2MASS
$K_{\rm s}$ (mag)	6.98 ± 0.03	2MASS
w_1 (mag)	6.91 ± 0.04	WISE
$w_2 \text{ (mag)} \dots \dots$	6.93 ± 0.02	WISE
w_3 (mag)	6.95 ± 0.02	WISE
$w_4 \pmod{\max(\max)}$	6.86 ± 0.05	WISE
	Spectroscopic Da	ata in TFOP
UT 2019-02-09	SNR=34	NRES@LCO
UT 2019-02-19	SNR=40	NRES@LCO
UT 2019-02-23	SNR=40	NRES@LCO
UT 2019-02-27	SNR=41	NRES@LCO
UT 2019-02-23	SNR=86	CHIRON@SMARTS
UT 2019-08-09	SNR=70	CHIRON@SMARTS
UT 2021-01-10	SNR=63	CHIRON@SMARTS
UT 2021-08-28	SNR=50	CHIRON@SMARTS

2.2. Stellar parameters

Light curves from photometric transits only allow the measurement of the planet-to-star radius ratio, hence we need to accurately determine the stellar radius in order to infer the radii of the transiting planets.

In the literature there are several estimates for the mass and radius of HD 22946: i) the values from Gaia DR2 (Gaia Collaboration et al. 2018), and ii) the values obtained by Kervella et al. (2019, hereinafter K19) by using the V, V - K surface brightness relations for the radius (Kervella et al. 2004) and the isochrone fitting method for the mass (Girardi et al. 2000). Both these approaches depend on the measured parallax. Besides these estimates, we can use two different and independent methods to infer the radius of the star and the other stellar parameters. The first method relies on the analysis of the spectra collected through the TESS Followup Observing Program

(TFOP⁴, (Collins 2019)): high-resolution, high signal-to-noise (SNR) optical spectra of HD 22946 were taken at 4 different epochs with both the Network of Robotic Echelle Spectrographs (NRES) (Siverd et al. 2018) of the Las Cumbres Observatory (LCO; Brown et al. 2013a) (Siverd et al. 2016) and the CHI-RON spectrograph (Tokovinin 2018a) on the 1.5m Small and Moderate Aperture Research Telescope System (SMARTS) at the Cerro Tololo Inter-American Observatory (CTIO).

The observations are detailed in Table 1. NRES data cover a 380-860 nm spectral band with a resolution of 53,000 and were processed by the dedicated data reduction pipeline (Mc-Cully et al. 2018a), and then analyzed by using SpecMatch (Petigura et al. 2017). CHIRON spectra cover the 450-890 nm range with a resolution of 80000. The spectra were collected by the instrument team and reduced following the optimal ex-

⁴ https://tess.mit.edu/followup/

traction method described in Paredes et al. (2021). We derived radial velocities by fitting the line profiles of the spectra, extracted via Least Squares Deconvolution (LSD) of the observed spectrum against synthetic templates (Donati et al. 1997), and we estimated the spectroscopic stellar parameters by using the gradient-boosting regressor implemented in the scikit-learn python module, previously trained on spectra classified by SPC (Buchhave et al. 2012). Our analysis of the line profiles reveals negligible rotational broadening and no evidence of double lines, suggesting that the star is inactive and not host to unresolved, close-in, stellar companions.

In Table 2 we list the radial velocity V_{rad} measured at each epoch, and the weighted average with associated uncertainties for all the stellar parameters obtained by the spectral analysis, with the prior of the Gaia parallax.

We note that the V_{rad} measured at the different epochs from the same spectrograph are all consistent with each other. A zeropoint discrepancy among the measures obtained from the two instruments is expected. NRES observations do not show significant variation at the level of 600 m/s over 18 days while CH-IRON spectra do not show significant variation at the level of 100 m/s over 2.5 years. Moreover, the V_{rad} from Gaia DR2 also shows a small uncertainty, i.e. a small *rms* from the different epoch observations, and is consistent with the value measured from the NRES spectra. All the above evidence further confirm that the star does not have a bound companion.

The second method used to obtain stellar parameters is based on the fitting of the spectral energy distribution (SED), i.e., the flux emitted by the star across a broad wavelength range, which yields a semi-empirical determination of the stellar radius, mass, and age.

We used the photometric data listed in Table 1, following the method described in Stassun & Torres (2016) and Stassun & Torres (2018). We retrieved public photometric data in different passbands as follows:

i) FUV and NUV magnitudes from the GALEX catalog; ii) *BV* magnitudes from the catalog of Mermilliod (2006); iii) $B_T V_T$ magnitudes from the Tycho-2 catalog; iv) $GG_{BP}G_{RP}$ magnitudes from Gaia EDR3; v) *JHK*_S magnitudes from the 2MASS catalog (Skrutskie et al. 2006; Cutri et al. 2003); and vi) magnitudes in the mid-infrared bands at 3.4, 4.6, 12 and 22 μ m from the WISE catalog (Wright et al. 2010). The full wavelength range covered by the data goes from 0.15 to 22 μ m, as shown in Figure 2.

We performed a fit using the Kurucz stellar atmosphere models, adopting the spectroscopically determined $T_{\rm eff}$, log g, and [Fe/H], as well as the $v \sin i$ from Pribulla et al. (2014). The remaining free parameter is the interstellar extinction, A_V , which we limited to the maximum line-of-sight value from the dust maps of Schlegel et al. (1998). The fit is reasonably good, with a reduced χ^2 of 1.9 and best fit $A_V = 0.02 \pm 0.01$. Integrating the (unreddened) SED gives $F_{bol}=1.289\pm0.030 \ 10^{-8}$ erg s⁻¹ cm⁻¹, that, combined with the Gaia EDR3 parallax (with no offset applied; see, e.g., Stassun & Torres 2021) gives $R_{\star} = 1.139 \pm 0.035 R_{\odot}$. With the spectroscopic log g, we then calculate a value of $M_{\star}=1.13\pm0.12M_{\odot}$, which is consistent with that determined by adopting the Torres et al. (2010) empirical relations, namely $M_{\star} = 1.12 \pm 0.06 M_{\odot}$. Finally, we can estimate the stellar rotation period via the observed R'_{HK} activity index (Murgas et al. 2013) and the empirical rotation-activity relations of Mamajek & Hillenbrand (2008), which yields $P_{\rm rot}$ = 10.6 ± 0.9 days . Using these rotation and activity measures together with the activity-age relations of Mamajek & Hillenbrand (2008) gives an estimated stellar age of $\tau_{\star} = 5 \pm 1$ Gyr.




Fig. 3. TESS image of HD 22946 taken during Sector 4 observations. The target is in the center, labeled with a "1" and a white cross. Nearby sources down to +10 mag are shown as smaller red dots and labeled in order of distance from the target. The SPOC-based, optimal aperture for the specific sector is overplotted in red. The color scheme represents the flux scale on the pixels. This image has been created using the tpfplotter python package (Aller et al. 2020).

The derived stellar parameters from both methods are consistent with each other and listed in Table 2, together with the best values adopted through the next steps of the analysis, that were obtained by performing the error-weighted average of all the available parameters.

Summarising, we found that HD 22946 is a late-type F main-sequence star, with $T_{\rm eff}$ =6040±48 K, log g=4.26±0.15 dex, [Fe/H]=-0.14 ± 0.07 dex, M_{\star} =1.104±0.012 M_{\odot} and R_{\star} = 1.157±0.025 R_{\odot} .

3. Discovery and characterization of HD 22946 planets

In this Section we present the TESS observations of HD 22946 and the analysis performed to detect and characterize the transit signals in its light curve.

3.1. TESS observations

TESS observed HD 22946 with Camera 3 in sectors 3 and 4 (2018-Sep-20 to 2018-Oct-18 and 2018-Oct-18 to 2018-Nov-15) during its primary mission and sectors 30 and 31 (2020-Sep-22 to 2020-Oct-21 and 2020-Oct-21 to 2020-Nov-19) during the extended mission. Fig.3 displays a single TESS frame of HD 22946 taken during Sector 4 observations. The first planet search has been carried out with the Science Processing Operation Center (SPOC) pipeline (Jenkins et al. 2016). The SPOC pipeline is designed to identify potential exoplanet transits using a wavelet-based, adaptive noise-compensating matched filter with the Transiting Planet Search code (Jenkins 2002). Subsequently, it fits a transit model to the detected signals (Li et al. 2019) and automatically performs a suite of tests to vet their planetary nature (Twicken et al. 2018). These are then collected in a Data Validation Report (DVR), available on both the

Table 2. Stellar data

Parameter	Value	Source
	Literature Data	
$T_{\rm eff}$ (K)	6115 ± 324	Gaia DR2
A_G (mag)	0.05 ± 0.5	Gaia DR2
$V_{\rm rad} ({\rm km \ s^{-1}})$	16.91 ± 0.15	Gaia DR2
$R_{\star}(R_{\odot})$	1.14 ± 0.11	Gaia DR2
$M_{\star} (M_{\odot})$	1.158 ± 0.058	K19
$R_{\star}(R_{\odot})$	1.115 ± 0.056	K19
$V_{\rm rad} ({\rm km \ s^{-1}})$	16.15 ± 0.04	Pribulla et al. (2014)
$v \sin i (\mathrm{km} \mathrm{s}^{-1})$	7.05 ± 0.2	Pribulla et al. (2014)
Spectral Type	F7/F8V	Pribulla et al. (2014)
	High-resolution spectroscopy	
$V_{\rm rad}~({\rm km~s^{-1}})$	16.6 ± 0.3	NRES@LCO
$V_{\rm rad} ({\rm km}{\rm s}^{-1})$	16.2 ± 0.3	NRES@LCO
$V_{\rm rad} ({\rm km}{\rm s}^{-1})$	16.5 ± 0.8	NRES@LCO
$V_{\rm rad} ({\rm km}{\rm s}^{-1})$	16.8 ± 0.4	NRES@LCO
$V_{\rm rad}$ (km s ⁻¹)	17.589 ± 0.033	CHIRON@SMARTS
$V_{\rm rad}$ (km s ⁻¹)	17.539 ± 0.033 17.624 ± 0.022	CHIRON@SMARTS CHIRON@SMARTS
$V_{\rm rad}$ (km s ⁻¹)		
	17.685 ± 0.038 17.620 ± 0.027	CHIRON@SMARTS
$V_{\rm rad}~({\rm km~s^{-1}})$	17.620 ± 0.037	CHIRON@SMARTS
$T_{\rm eff}$ (K)	$6210 \pm {}^{173}_{158}$	This work, NRES
$\log g$	4.4 ± 0.2	This work, NRES
[Fe/H]	$-0.05 \pm ^{0.08}_{-0.05}$	This work, NRES
$v \sin i (\mathrm{km} \mathrm{s}^{-1})$	3.4 ± 0.9	This work, NRES
M_{\star}	1.104 ± 0.012	This work, NRES
R_{\star}	1.157 ± 0.025	This work, NRES
$T_{\rm eff}$ (K)	6026 ± 50	This work, CHIRON
$\log g$	4.22 ± 0.10	This work, CHIRON
[Fe/H]	-0.22 ± 0.08	This work, CHIRON
$v\sin i$ (km s ⁻¹)	3.2 ± 0.5	This work, CHIRON
	SED fitting from broad-band photometry	
A_V	0.02 ± -0.01	This work
$F_{bol} (\text{erg s}^{-1} \text{ cm}^{-2})$	1.289 ± 0.030	This work
$R_{\star} (R_{\odot})$	1.139 ± 0.035	This work
M_{\star} (M_{\odot})	1.12 ± 0.06	from $\log g$ and R_{\star}
$M_{\star}~(M_{\odot})$	1.13 ± 0.12	from Torres et al. (2010)'s empirical relations
Age (Gyr)	5 ± 1	This work
	Best values adopted through this work	
$T_{\rm eff}$ (K)	6040 ± 48	This work
$\log g$	4.26 ± 0.15	This work
[Fe/H]	-0.14 ± 0.07	This work
$R_{\star} (R_{\odot})$	1.157 ± 0.025	This work
$M_{\star}~(M_{\odot})$	1.104 ± 0.012	This work

ExoFOP-TESS and MAST⁵ archives. At this point, the pipeline masks the detected signals and runs from the top to identify new signatures. It stops after eight tries or in case the latest found signal is below a certain SNR threshold. Finally, the pipeline generates two different light curves for the detected TOIs: the first one is extracted through simple-aperture photometry (SAP-FLUX; Twicken et al. (2010); Morris et al. (2020)), whilst the other one has an additional pre-search data conditioning procedure (PDCSAP-FLUX; Stumpe et al. (2014)). This final step re-

⁵ https://mast.stsci.edu/

moves long-term trends and contamination due to nearby stars. Two planet candidates have been discovered during this preliminary analysis and added as TESS TOI-411.01 and 411.02 to ExoFOP-TESS. These candidates have periods of 9.57 days and 4.04 days, with transit epochs at 1385.72367 and 1386.189174 Barycentric TESS Julian Date (BTJD⁶), respectively. The full TESS light curve is shown in Fig. 4. The transits of the candidates are highlighted.

Article number, page 5 of 21

 $^{^{6}}$ BTJD = BJD - 2457000.0




Fig. 4. Full light curve of TESS sectors 3, 4 (top panel) and 30, 31 (bottom panel). The blue line is the best fit model. The transits of planet b and c are highlighted with orange and green dots, respectively. The single transit event is highlighted with a red dot.

3.2. Light curve analysis

We downloaded and worked on the PDCSAP light curve for TOI 411 using the lightkurve package (Lightkurve Collaboration et al. 2018). This code makes use of an automatically-generated optimal aperture but we checked that different tentative choices of apertures around this target do not affect the depth of the transit signals. Starting with this time series, we first recover the candidate events and subsequently fit them to retrieve physical and orbital parameters.

3.3. Candidates detection

We used the Transit Least Squares (TLS) package (Hippke & Heller 2019) to recover the candidate signals. This code allows searching for periodic, transit-like signals in a light curve by brute-force fitting templates of transits built over a grid of parameters. We decided to adopt this code with respect to the more commonly used Box Least Squares (BLS, Kovács et al. (2002)) algorithm, because it also accounts for the effect of stellar limbdarkening on the shape of transits, and, according to the authors, this improvement yields a $\sim 10\%$ higher detection efficiency (and similarly lower false alarm rates) compared to BLS, which is especially relevant for low SNR signals. The brute-force fitting consists of the phase-folding of the light curve over a grid of trial periods, transit epochs and duration, and computing the χ^2 at each solution. The solution is then found by minimizing the χ^2 . The period grid is a uniformly-spaced set of N possible values within a given range. The upper limit of the range is set by the minimum number of transits we want to find over the entire observation window, which we set to two. The lower limit is such that the candidate orbits just wide of the Roche limit of its star (in the assumption of a planet density of $\rho_p = 1 \text{ g cm}^{-3}$). In the case of HD 22946, this limit corresponds to 0.6 days. We performed the first TLS run independently on the couples of sectors [3,4] and [30,31] to ease the computational burden of the search. The first run on sectors 3 and 4 yields a periodic event with a period of ~ 9.57 days, which is consistent with the ExoFOP TESS listed period for TOI 411.01. Masking this signal and running a second search, we recover the events associated with TOI 411.02 on a \sim 4.04 days period. A third TLS run does not return consistent periodic events. The same results are found for sectors 30 and 31. In addition, by visually inspecting the available sectors,

we notice two additional events: a transit-like feature in sector 4 at BTJD \sim 1425 and another dip in sector 30 at BTJD \sim 2136.5. However, the latter event presents an asymmetric shape and it overlaps the momentum dump reported in TESS Data Release Notes no.45 (DRN45, and Fig.4 and Fig.7 therein) on MAST⁷ at the same epoch. The former event, instead, does not correspond to any reported spacecraft maneuver epoch. Hence, we only consider the sector 4 transit-like signal hereafter. This newly found signal is then a single event candidate exoplanet whose period is yet to be defined (see Fig. 4 of this work).

This single event candidate was also recently reported independently on ExoFOP-TESS (as TIC 100990000.03) by A.S., a member of the Planet Hunters TESS citizen science project (Eisner et al. 2021).

We note that TESS is not scheduled to observe this target again during its upcoming extended mission and therefore follow-up observations with other facilities will be needed to confirm this newly found candidate exoplanet.

3.4. Transit modeling

We used exoplanet, a code for probabilistic modeling of exoplanet transits (Foreman-Mackey et al. 2021), to measure the properties of the transiting planets. The model we set up consisted of four elements: three planet transit components with Keplerian orbits and limb-darkened transits, and a Gaussian Process (GP) component that models residual stellar variability. We combined the data from all the sectors into one time series with the same median normalized brightness. The planet models were computed with exoplanet using STARRY (Luger et al. 2019), while the GP was computed using celerite (Foreman-Mackey et al. 2017; Foreman-Mackey 2018). The GP component is described as a stochastically-driven, damped harmonic oscillator with parameters of log(S_0) and log(ω_0), where the power spectrum of the GP is

$$S(\omega) = \sqrt{\frac{2}{\pi}} \frac{S_0 \,\omega_0^4}{(\omega^2 - \omega_0^2)^2 + \omega^2 \,\omega_0^2/Q^2},\tag{1}$$

and a white noise term, with a model parameter of the log variance. We fixed Q to 1/3 and put wide Gaussian priors on $\log(S_0)$

⁷ https://archive.stsci.edu/tess/tess_drn.html

and $log(\omega_0)$ with means of the log variance, and one log of onetenth of a cycle, respectively, and a standard deviation on the priors of 10. This form of GP was chosen because it enables us to model a wide range of low-frequency astrophysical and instrumental signals without requiring a physical model for the observed variability. The white noise term carried the same prior as $\log(S_0)$. The planet model was parameterized in terms of limb darkening, log stellar density and stellar radius for the three planets. Each individual planet was parameterized in terms of log orbital period, time of transit, planet-to-star radius ratio, impact parameter, orbital eccentricity and periastron angle at the time of transit. The time of transit was set to a transit near the center of the time-series to minimize correlations between transit epoch and orbital period. The stellar radius had a Gaussian prior of $1.157 \pm 0.025 R_{\odot}$. The log mean stellar density, in cgs units, had a Gaussian prior with a mean of log 1.1 and standard deviation of 0.18 dex. The limb darkening followed the Kipping (2013a) parameterization. The log orbital periods, time of transit, and log planet-to-star radius ratio of the three candidates had Gaussian priors with means at the values listed on ExoFOP-TESS (aside from the single transiting planet where we fixed the orbital period). The impact parameter had a uniform prior between zero and one plus the planet-to-star radius ratio. Eccentricity had a beta prior with $\alpha = 0.867$ and $\beta = 3.03$ (as suggested by Kipping 2013b), and was bounded between zero and one. The periastron angle at transit was sampled in vector space to avoid the sampler seeing a discontinuity at values of π . We sampled the posterior distribution of the model parameters using the No U-turn Sampler (NUTS, Hoffman & Gelman 2014) which is a form of Hamiltonian Monte Carlo, as implemented in PyMC3 (Salvatier et al. 2016). We ran 4 simultaneous chains, with 3000 tuning steps, and 2000 draws in the final sample. The phasefolded transits of the three candidates, along with best-fitting transit models, are shown in Figure 5, and the model parameters are provided in Table 3.5. Note that, by using the transit duration and depth along with stellar mass and radius, we can estimate the period through Kepler's third law in the assumption of planar and circular orbit, as shown in Section 6.5 of Seager & Mallén-Ornelas (2003). This yields a period of approximately 46 ± 4 days. A period between 43 and 46 days would also be consistent with the possible overlap of a secondary transit corresponding to momentum dump in sector 30, with a missing signal that would fall in the downlink gap of sector 31 and just short of the beginning of observations for sector 3. Moreover, if we assume that the signal at BJD at 2136.5 is a secondary transit happening at the same time of the reported momentum dump in sector 30, we find a period of 44.47 days, which is also consistent with our rough estimate. On the basis of TESS data only, we are not able to establish if the signal that we observed in sector 30 at BJD at 2136.5 is only due to an instrumental artifact or if it also includes the transit of planet d.

3.5. Additional radial velocity data

By searching the ESO archive, we found that TOI-411 was observed by the Echelle SPectrograph for Rocky Exoplanets and Stable Spectroscopic Observations (ESPRESSO; Pepe et al. (2014)) between the 10th of February and the 17th of March 2019 (program ID: 0102.C-0456; PI: Van Eylen). ESPRESSO is a high-resolution (\mathbb{R} ~140,000) spectrograph operating at the VLT at ESO's Paranal Observatory, capable of measuring radial velocities of bright stars (V< 8 mag) with a precision of 10 *cm/s* in the 377–790 nm range. TOI-411 was observed 14 times in roughly one month, with typical exposure of 600s and SNR




Fig. 5. Folded TESS light curves from sectors 3, 4, 30 and 31 corresponding to the transits of planet b (top panel), planet c (middle panel) and the single event transit of candidate planet d from sector 4 (bottom panel). The exoplanet fits of the transits are overplotted (colored lines) while blue points are binned data points.

ranging from 120 to 243. The observations are summarised in Table3.5. We looked at the radial velocities publicly available in the archive, which are derived from the cross-correlation of the stellar spectra with a G2 template. The data is shown in Fig.6.

We modelled the ESPRESSO radial velocity data using exoplanet (Foreman-Mackey et al. 2017; Foreman-Mackey 2018).




Fig. 6. Upper panel: ESPRESSO precise radial velocity measurements of TOI-411, taken over February and March 2019. Lower panel: the pRVs have been phase-folded on the periods of planet candidates TOI-411 b and c.

Table 3. ESPRESSO radial velocity measurements of HD 22946

$CCF_{RV}[km/s]$	$CCF_{RV_{err}}[km/s]$	$\langle SNR \rangle$	MJD
16.864287	0.000561	181	58524.05723591
16.865981	0.000627	164	58525.05175402
16.868350	0.000553	184	58526.09212435
16.866658	0.001012	109	58527.12891579
16.865454	0.000619	168	58535.12063523
16.864550	0.000496	203	58540.03340549
16.869616	0.000916	119	58550.06931525
16.866772	0.000763	139	58552.06584297
16.876263	0.000540	188	58553.01522444
16.868946	0.000662	156	58555.99860106
16.870850	0.000447	224	58557.00297657
16.870675	0.000487	208	58557.06290428
16.870844	0.000482	210	58558.02527738
16.870741	0.000766	138	58559.03833688

Specifically, we included the V_rad data with the transit model and performed a joint, self-consistent model. As parts of the V rad component of the model, we included a periodic Gaussian Process with a characteristic timescale of 50 days and an amplitude of 2 m/s. The values chose were selected to be longer than any Keplerian signal from the planets, and designed to encompass any long-term variations in the data. The period of planet d was fixed to 46 days. The posterior distributions of the semiamplitudes are consistent with non-detection of any signal in the radial velocity data, with 3σ upper limits of 4.1, 3.5, 8.4 m/s for planets b, c and d, respectively. These values for the semiamplitudes yield upper mass limits of ~ 11, 14.5 and 24.5 M_{\oplus} . The masses are forecast to be in the region of 3.6, 8.0, and 11 M_{\oplus} (Chen & Kipping 2017), and therefore the non-detection of any mass signal is not unexpected. Finally, there is evidence for a slope in the RV data, although the significance is not high. In the month that the RV data cover, the star shows show a 6 m/s change in measured radial velocity. This signal is consistent with an amplitude that might be seen if the source of the signal were to be stellar rotation or RV jitter (Luhn et al. 2020). If the source is not stellar or instrumental noise, then the slope could be due to an external body such as another planet or stellar companion. However, the available data is not sufficient to make predictions beyond saying that the data is not inconsistent with a companion more massive than Neptune on an orbital period longer than about 30 days.

Article number, page 8 of 21

4. Ruling out false positive scenarios

TESS is designed to have a large field of view $(24^{\circ} \times 96^{\circ})$ to survey 85% of the entire sky within the two years of the nominal mission. As a consequence, it also has a large pixel scale (21" per pixel), with a focus-limited PSF that can be as large as 1'. This implies that the collected time-series photometry of a given target can be contaminated by nearby or background/foreground unresolved sources. Specifically, the aperture used to extract the light curve could contain more sources. The light contribution of this other source could dilute the transit signal, resulting in an underestimation of the associated planetary radius. Or the additional source could be an eclipsing binary of which we can only observe one set of eclipses due to signal dilution caused by the light collected from the target star, producing a false positive. Therefore, the various scenarios we have to rule out include: a) possible instrumental (systematic) effects; b) the host star being in an eclipsing binary or multiple system; c) the host star being close to or aligned with a foreground or background star eclipsed by a stellar companion or transited by a planet.

To address the first scenario, we inspected TESS Data Releases for the interested sectors to look for instrument malfunctioning and reported artifacts. As stated in Section 3.3, we found a feature at 2136.5 BTJD associated with one of the momentum dumps, i.e. times at which the reaction wheel speeds of the satellite were reset. We therefore decided to discard this event as an instrumental artifact. We did not find any other feature of similar nature in the entire light curve.

To clear HD 22946 from the remaining false positive scenarios, we first directly analyze *TESS* images and light curves, by using the DAVE pipeline (Kostov et al. 2019), as described in subsection 4.1.

Secondly, we investigate the veracity of the transit-like signals using additional ground-based time-series and highresolution imaging collected within TFOP, as described in subsections 4.3 and 4.2. All of these tests have exclusively been performed on planet candidates b and c, since they are based on the detection of multiple events and therefore scheduled around the predictions made from the known periods. However, we note that if we assume a population of false positive randomly distributed over the entire sky, a star with at least one (confirmed) transiting planet is more likely to show signals of a second transiting planet rather than being a false positive (Latham et al. 2011; Lissauer et al. 2012; Guerrero & TESS Science Office 2021). This multiplicity factor increases the probability that a planet candidate is a true planet rather than a false positive by a factor of 20-50 (Rowe et al. 2014; Lissauer et al. 2014; Guerrero & TESS Science Office 2021). These considerations help us to increase our confidence in the planetary nature of all three candidates investigated in this work.

To quantify this confidence, we computed the Bayesian probability for the signals being false positives with triceratops (Giacalone & Dressing 2020) and vespa (Morton 2015) in subsection 4.4.

4.1. DAVE analysis

In order to tackle the false-positive scenarios b) and c), we first use the DAVE package, following the workflow of Cacciapuoti et al. (2022). We use different modules of the code, with the following rationale.

centroids generates the in-transit and the two (before and after the event) out-of-transit images for each transit. Then, it

Table 4. Planet parameters. Th	priors for each fitted quantity an	re explained in the text (Section 3.4).
--------------------------------	------------------------------------	---

Parameter	Median	$+1\sigma$	-1 <i>o</i>
<u>S4</u>	Model Parameters		
Star	1 10	0.10	0.16
$\rho [\text{g cm}^{-3}]$	1.10	0.19	0.16
Limb darkening u_1	0.18	0.20	0.13
Limb darkening u_2	0.19	0.28	0.24
TOI-411 b			
<i>T</i> ⁰ (BJD - 2457000)	1385.729875	0.004361	0.00119
ln Period [days]	1.396319	0.000006	0.00001
Impact parameter	0.29	0.23	0.12
Depth (ppm)	201	17	16
eccentricity	0.126	0.190	0.009
ω [radians]	-0.7	2.2	2.9
TOI-411 c			
<i>T</i> ⁰ (BJD - 2457000)	1386.188119	0.001278	0.00131
ln Period [days]	2.2589566	0.0000027	0.000002
Impact parameter	0.42	0.20	0.27
Depth (ppm)	502	32	30
eccentricity	0.16	0.17	0.10
ω [radians]	0.2	2.2	2.4
Single Transit Event T_0 (BJD - 2457000)	1425.164717	0.002288	0.00190
$\ln Period$ [days]	1723.107/1/	0.002288	0.00190
Impact parameter	0.35	0.21	0.23
Depth (ppm)	704	64	65
eccentricity	-	04	05
ω [radians]	-	-	-
w [laulans]	-	-	-
TOI-411 b	Derived Parameters		
	4.040301	0.000023	0.00004
Period [days]	0.01358	0.000023	0.00004
R_p/R_*	1.72	0.00000	0.0005
Radius $[R_{\oplus}]$			
a/R_*	9.84	0.53	0.51
a [AU]	0.0528	0.0030	0.0030
Inclination (deg)	88.3	1.1	1.2
Transit duration (hours)	2.96	0.38	0.53
Equilibrium temperature (K)	1378	36	36
TOI-411 c	0.770000	0.0000	0.00000
Period [days]	9.573096	0.000026	0.00002
R_p/R_*	0.02161	0.00077	0.00075
	2.74	0.14	0.14
Radius $[R_{\oplus}]$			
a/R_*	17.49	0.95	0.91
<i>a/R</i> _* <i>a</i> [AU]	17.49 0.0939		0.0054
a/R_* a [AU] Inclination (deg)	17.49	0.95	
a/R_* a [AU] Inclination (deg) Transit duration (hours)	17.49 0.0939	0.95 0.0054	0.0054
a/R_* a [AU] Inclination (deg)	17.49 0.0939 88.57	0.95 0.0054 0.86	0.0054 0.53
a/R_* a [AU] Inclination (deg) Transit duration (hours)	17.49 0.0939 88.57 3.95	$\begin{array}{c} 0.95 \\ 0.0054 \\ 0.86 \\ 0.82 \end{array}$	0.0054 0.53 0.80
a/R_* a [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K)	17.49 0.0939 88.57 3.95	$\begin{array}{c} 0.95 \\ 0.0054 \\ 0.86 \\ 0.82 \end{array}$	0.0054 0.53 0.80
<i>a/R</i> _* <i>a</i> [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K) Single Transit Event Period [days]	17.49 0.0939 88.57 3.95	$\begin{array}{c} 0.95 \\ 0.0054 \\ 0.86 \\ 0.82 \end{array}$	0.0054 0.53 0.80
a/R_* a [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K) Single Transit Event	17.49 0.0939 88.57 3.95 1033	0.95 0.0054 0.86 0.82 27	0.0054 0.53 0.80 27
a/R_* a [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K) Single Transit Event Period [days] R_p/R_* Radius $[R_{\oplus}]$	17.49 0.0939 88.57 3.95 1033	0.95 0.0054 0.86 0.82 27 0.0012	0.0054 0.53 0.80 27
a/R_* a [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K) Single Transit Event Period [days] R_p/R_* Radius $[R_{\oplus}]$ a/R_*	17.49 0.0939 88.57 3.95 1033	0.95 0.0054 0.86 0.82 27 0.0012 0.19	0.0054 0.53 0.80 27 .0012 0.19
a/R_* a [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K) Single Transit Event Period [days] R_p/R_* Radius $[R_{\oplus}]$ a/R_* a [AU]	17.49 0.0939 88.57 3.95 1033	0.95 0.0054 0.86 0.82 27 0.0012 0.19	0.0054 0.53 0.80 27 .0012 0.19
a/R_* a [AU] Inclination (deg) Transit duration (hours) Equilibrium temperature (K) Single Transit Event Period [days] R_p/R_* Radius $[R_{\oplus}]$ a/R_*	17.49 0.0939 88.57 3.95 1033	0.95 0.0054 0.86 0.82 27 0.0012 0.19	0.0054 0.53 0.80 27 .0012 0.19

Article number, page 9 of 21

subtracts the overall in-transit image from the overall out-oftransit image to produce a difference image. Finally, it measures the center-of-light for each difference image by fitting the *TESS* Pixel Response Function (PRF⁸)to the image and computes the overall photocenter by averaging over all the events examined at the previous step. The aim of this procedure is to pinpoint the true source of the transit-like events. See Kostov et al. (2019) for further details.

- modelshift phase-folds the light curve and convolves it with the best-fit trapezoid transit model, thus highlighting light curve features. Furthermore, it shows the average of the input primary signals, the average of the odd and even signals, the most prominent secondary, tertiary and positive features. These results allow us to investigate the possibility that the source of the signal is an eclipsing binary. See Kostov et al. (2019) for further details.
- DAVE finally runs the astropy-implemented Lomb-Scargle periodogram and generates a PDF file showing both the light curve phase-folded on the period of the inspected signal and the LS period. This test is performed to check whether light curve modulations occur on (half) the orbital period of the candidate. This is typical of beaming, reflection and ellipsoidal effects of binary systems (see e.g., Morris & Naftilan 1993; Faigler & Mazeh 2011; Shporer 2017).

DAVE results for both TOI 411.01 and TOI 411.02 show no significant additional eclipse in the light curves, no odd-even differences in consecutive signals, nor flag the shapes of the transits. Even though some transits of planet b and c produce unreliable photocenter difference images due to their low signal-to-noise ratio, the ones that are reliable confirm that the target star is the true source of the signal. For every sector, the average of all photocenters for each difference image of planets b and c falls on the target star. Since no significant shift is found, HD 22946 can be considered the source of the recovered signals, at TESS resolution limit. Regarding planet candidate d, no reliable photocenter measurement can be drawn due to the fact that only a single event has been observed by *TESS*. Modelshift also requires periodic signals, so it cannot be applied to the single-event candidate d.

Several resulting images of the centroids module and examples of Modelshift's results are shown in the Appendix (Fig. A.1, Fig. A.2, Fig.A.3 and Fig.A.4).

Finally, it is worth mentioning that TOI-411 *b* passed all the automated SPOC pipeline Data Validation (DV) diagnostic tests but the difference-image centroiding test for sectors 4 and 31. However, the difference-image centroiding tests for this planet are affected by the fact that TOI-411 appears to be slightly saturated and the SNR of the transit signal is low (\sim 10). TOI-411 *c* passed all the DV diagnostic tests, including the difference-image centroiding tests for transit signatures passed the odd/event transit depth tests reported by the DV. The summary of these results can be found on ExoFOP-TESS⁹.

4.2. Ruling out nearby eclipsing binaries

We acquired ground-based time-series follow-up photometry of the stars in the field around TOI-411 as part of the *TESS* Follow-




Fig. 7. Las Cumbres Observatory image of the host star HD 22946 in the rp filter. The 10 sources identified by pink circles as T2 through T11 are within a 2.5' radius of TOI 411 and are potentially bright enough (see text) to cause the TOI 411 b/c signals. The nine closest nearby sources are also displayed in Fig.3 and fall just outside the automatic, SPOC-based aperture used in Sector 4.

up Observing Program (TFOP; Collins 2019)¹⁰. Observations were scheduled to cover the times of transit of TOI-411.01 and TOI-411.02, predicted by the initial published Quick Look Pipeline (QLP, (Huang et al. 2020)) ephemerides from *TESS* sectors 3 and 4 using the TESS Transit Finder, which is a customized version of the Tapir software package (Jensen 2013). If the events detected in the *TESS* data are indeed on-target, the shallow QLP reported depths of 280 ppm and 166 ppm for TOI-411.01 and TOI-411.02, respectively, would not generally be detectable in ground-based observations. Instead, we saturated the bright star TOI-411 to enable the extraction of light curves of nearby fainter stars to attempt to rule out or identify nearby eclipsing binaries (NEBs) as potential sources of *TESS* detection.

We observed an ingress plus about 50% of a predicted TOI-411.01 event and a full transit of TOI-411.02 using the Las Cumbres Observatory Global Telescope (LCOGT; Brown et al. 2013b) 1.0 m network nodes at the South Africa Astronomical Observatory and Cerro Tololo Inter-American Observatory, respectively. The TOI-411.01 observation was on UTC 2019 February 24 in Sloan r' band and the TOI-411.02 observation was on UTC 2020 December 02 in Sloan i' band. The 1 m telescopes are equipped with 4096 × 4096 SINISTRO cameras having an image scale of 0".389 per pixel, resulting in a $26' \times 26'$ field of view. The images were calibrated by the standard LCOGT BANZAI pipeline (McCully et al. 2018b). Photometric data were extracted using AstroImageJ (Collins et al. 2017).

Additionally, full transit duration observations were performed with the 0.305m Perth Exoplanet Survey Telescope (PEST) in the Rc filter on the nights of 2019-02-16 (TOI-411.02) and 2020-12-06 (TOI-411.01). The data reduction and the aper-

⁸ For more information about TESS PRF see https: //archive.stsci.edu/files/live/sites/mast/files/home/ missions-and-data/active-missions/tess/_documents/ TESS_Instrument_Handbook_v0.1.pdf

⁹ https://exofop.ipac.caltech.edu/tess/target.php?id=100990000

¹⁰ https://tess.mit.edu/followup

ture photometry were performed by using a custom pipeline based on C-Munipack¹¹.

Although the original analyses were based on the initial published OLP ephemerides from TESS sectors 3 and 4, we reanalyzed the data relative to the TOI-411.01 and TOI-411.02 ephemerides derived in this work. We checked for possible NEBs that could be contaminating the SPOC and QLP photometric apertures, which generally extend $\sim 1'$ from the target star. To account for possible contamination from the wings of neighboring star PSFs, we searched for NEBs in all known Gaia EDR3 and TIC version 8 nearby stars out to 2.5 from TOI-411 that are possibly bright enough in TESS band to produce the TESS detection (assuming a 100% eclipse and 100% contamination of the TESS aperture). In order to account for possible delta-magnitude differences between TESS-band and the followup filter bands, we checked stars that are an extra 0.5 magnitudes fainter in TESS-band than needed. We consider a star cleared of an NEB if the RMS of its 10-minute binned light curve is more than a factor of 5 smaller than the adjusted expected NEB depth in the star (adjusted to allow for the potential TESS-band deltamagnitude difference). We then visually inspect each neighboring star's light curve to ensure no obvious eclipse-like signal.

The transit depths we derive in this work are deeper than the initial QLP depths. To be conservative when checking for potential NEBs, we used the shallower TOI-411.02 QLP depth of 166 ppm to calculate adjusted expected depths in all of the nearby stars. This resulted in a check of the 10 stars labeled T2 through T11 in Figure 7. To simplify the presentation of the results, we also checked the same 10 stars for the deeper transit of TOI-411.01. The faintest star we checked is T2 (TIC 100990001), which has a delta-TESS-band magnitude of 9.86 according to TIC version 8. Assuming its true delta magnitude is 9.36 in the follow-up filter bands, a 92% eclipse in T2 would produce a 166 ppm depth in TOI-411. We find that the RMS of all 10 light curves for each planet candidate are more than a factor of 5 smaller than the adjusted expected NEB depth in the respective star. All of our follow-up light curves and supporting results are available on the EXOFOP-TESS website¹².

Finally, we note that no observations could be carried out to perform a similar analysis for planet candidate d, since the orbital period is unknown and hence we cannot make reliable predictions for the epoch of the next transits.

4.3. Ruling out unresolved stellar companions

Unresolved, dimmer stars could fall in the same 21"-wide TESS pixel. If an additional source is indeed present, and if this source is an eclipsing binary, its (diluted) eclipses might be the signals we observe in the light curve. Besides, the light contamination due to this unresolved star (system) might dilute the transit candidates, resulting in underestimated radii for the exoplanets (see e.g., Furlan et al. 2017; Ciardi et al. 2015). To rule out stellar companions and foreground/background stars at close separations unresolved in TESS images, we obtained speckle imaging observations with the 8m Gemini-South telescope's Zorro instrument (Scott et al. 2021), which provides speckle imaging simultaneously in two bands centered at 562 nm and 832 nm, resulting in reconstructed images and robust contrast limits on companion detections (see Howell et al. 2011, 2016). TOI 411 was also observed with the 4.1m SOuthern Astrophysical Research (SOAR) telescope (Ziegler et al. 2020, 2021) speckle imaging




Fig. 8. Gemini-South Zorro's speckle high-contrast imaging observations of TOI-411 on UT 2020-11-25: 5σ sensitivity curves and reconstructed images. The data show that there are no close-in companions detected down to 5 magnitude fainter than our target at a separation of 0.1 arcseconds.

camera HRCam (Tokovinin 2018b). Specifically, HD 22946 was observed with Zorro on UT 2020-11-21 and 2020-11-25 in both bands. The estimated PSF at the time of observations was 0.02". The 5σ sensitivity curves and the reconstructed images for the two bands are shown in Figure 8. We find that TOI-411 has no companion brighter than about 5 to 8 magnitudes, respectively, from about 0.1" out to 1.2". Observations with the HRCam on the SOAR were carried out on the UT 2019-02-18 in the I-band, centered at 879nm. The estimated PSF at the epoch of the observations was 0.06". The auto-correlation functions for the data obtained with SOAR observations is shown in Fig.9. No nearby star is detected within 3" of TOI-411, which corresponds to 188 AU at the distance of the target. Finally, we note that the host star HD 22946 was also analyzed by Kervella et al. (2019, 2022) for proper motion anomalies that could be induced by an unresolved orbiting companion, resulting in a non-detection. The Gaia Renormalised Unit Weight Error (RUWE) of the star is indeed 1.039, a typical value of single bright stars (Gaia Collaboration et al. 2021; Ziegler et al. 2020), thus further confirming the single-star scenario.

4.4. Quantifying the false positive probability for different scenarios

To quantify the false positive probability (FPP) for this system, we use the triceratops (Giacalone et al. 2021; Giacalone & Dressing 2020) and vespa (Morton 2015) packages. triceratops is an algorithm that rules out astrophysical false positives by calculating and comparing the probabilities of various transit-producing scenarios. triceratops is specifically designed for *TESS* observations that consider transit scenarios originating from the target star, sources unresolved with the target star, and known nearby stars within 2.5 arcminutes from the target star. This tool encapsulates the total probability that a planet candidate is a false positive in the false positive probability (FPP) and the probability that the planet candidate is a false positive probability (NFPP). For a planet candidate to be validated,

¹¹ http://c-munipack.sourceforge.net/

¹² https://exofop.ipac.caltech.edu/tess




Fig. 9. Analysis of the SOARCam high contrast image of TOI 411: the two-dimensional autocorrelation function and the reconstructed image of the field are shown. The data show that there are no close-in companions detected within 3 arcseconds of TOI 411, which would show up as additional peaks in the autocorrelation function.

it must achieve FPP < 0.01 and NFPP < 0.001 (see Giacalone et al. 2021, for more details). As an additional constraint in our calculations, we fold in the speckle imaging follow-up observations discussed in the previous Section. Because these observations reveal no previously unresolved companions within their detection limits, incorporating the follow-up reduces the calculated probability of the transit originating from a bound or chance-aligned star within the resolution limits of the target star, thereby reducing the FPP of the planet candidate. We note that, since the true period of planet *d* is unknown, we cannot perform this kind of analysis for it.

Otherwise, we ran triceratops for both planet candidates b and c twenty times and calculated the mean and standard deviation of the resulting FPPs and NFPPs. We found FPP = $(0.88 \pm 1.28) \times 10^{-7}$ and FPP = $(3.56 \pm 1.27) \times 10^{-4}$ for TOI-411.01 and TOI-411.02, respectively. Because triceratops determines that no nearby stars are capable of being sources of astrophysical false positives, we find NFPP = 0 for both candidates. Based on these results, we consider planets b and c to be validated.

Additionally, we used vespa to independently validate the same signals. vespa compares transit signals to a number of false-positive scenarios including an unblended eclipsing binary (EB), a blended background EB, a hierarchical companion EB, and the 'double-period' EB scenario. It accounts for the period, depth, duration, and shape of each signal as well as the colors of the target star, spectroscopic and imaging follow-up observations, and simulations of the population and distribution of field stars and binary stars at the target's position. We ran vespa on the TESS light curves to calculate the FPP for each individual planetary signal after masking additional potential transits for each target signal. The observational constraints imposed by Gemini-South Zorro's high contrast imaging of TOI 411 were included in this case as well. vespa is particularly sensitive to the constraint on the maximum depth of potential secondary eclipses and the maximum radius at which vespa considers the influence of background binary systems. We used the DAVE analysis to estimate the secondary depth constraint and set the maximum radius parameter at 21" to emulate the size of a *TESS* pixel. Using these inputs, we calculated an FPP of 7.77×10^{-12} and 4.51×10^{-5} for TOI 411.01 and TOI 411.02, respectively. These values are well below the commonly suggested 0.01 threshold required to statistically validate these candidates as planets, in agreement with the results from triceratops.

5. The HD 22946 planetary system

In this section, we present the main properties of the planetary system orbiting HD 22946. The three planets are a $1.72\pm0.10 R_{\oplus}$ Super-Earth (planet *b*), a $2.74\pm0.14 R_{\oplus}$ (planet *c*) and a $3.23\pm0.19 R_{\oplus}$ (planet *d*) sub-Neptunes. In subsection 5.1, we estimate the planet masses and the expected semi-amplitudes of radial velocity curves that could be measured.

In subsection 5.3, we perform N-body simulations to assess the dynamical stability of the system, while in subsection 5.4 we investigate the dynamical packing of the system to determine whether additional planets might be orbiting HD 22956. Finally, we evaluate the capability of spectroscopic follow-ups with the JWST NIRISS instrument in subsection 5.5 for the detection of possible atmospheres.

5.1. Planet mass estimates

While TOI-411 has been observed with ESPRESSO and a total of 14 radial velocity measurements are available as explained in Section 3.4, we note that a significantly larger number of spectra would be needed to place meaningful mass constraints on the planets of the TOI-411 system. Given the lack of them, we use empirical radius-mass relations to estimate the planets' masses.

We make use of two different equations for the inner planet and the outer ones. In fact, while the well-known Chen & Kipping (2017) relations have been widely used in the literature to estimate the masses of exoplanets of any size, Otegi et al. (2020) showed that a slightly different equations better constrains Super-Earths ($1.5 < R/R_{\oplus} < 2$). Thus, adopting the planet radii in Table3.5 and using Chen & Kipping (2017) relations, we estimate the masses of planets *c* and *d* to be $M_c = 7.96 \pm 0.69M_{\oplus}$ and $M_d = 10.53 \pm 1.05M_{\oplus}$, respectively. On the other hand, considering the corrections of Otegi et al. (2020), we estimate a mass of $M_b = 6.29 \pm 1.30M_{\oplus}$ for planet *b*.

Given the brightness of the host star ($V \approx 8.3$ mag), these planets are optimal targets for mass measurements by means of radial-velocity curves. In fact, the computed mean masses yield $V_{\rm rad}$ semi-amplitudes of 2.36 ± 0.48 m/s, 2.24 ± 0.19 m/s and 1.76 ± 0.18 m/s, respectively, which are well within the capabilities of current Southern Hemisphere instruments such as HARPS (Pepe et al. 2002) and ESPRESSO (Pepe et al. 2014), assuming an intensive monitoring.

5.2. Transit timing variation analysis

In order to obtain independent estimates of the planetary masses, we performed a transit timing variation (TTVs) analysis, which is also able to find possible undetected planets. TOI-411 system was modeled with the TRANSITFIT5 transit modeling software (Rowe & Thompson 2015; Rowe 2016). We computed a multiplanet fit that includes all 3 planets and assumes non-interacting Keplerian orbits. The best-fit model was then used to create a theoretical light curve with 2 of the other planets removed, thus isolating a single planet. The single-planet light curve was then used to measure the center of transit times for each transit event.




Fig. 10. One of the simulated eccentricity evolution of the HD 22946 system over 10^5 years. In the case shown here, we assumed the following extreme initial conditions for planet *d*: initial semi-major axis of 0.19 AU and eccentricity of 0.4.

We did not find any obvious sign of TTVs. The period ratio of planets b and c are not close to strong orbital resonance and the potentially long period of d suggests all three planets are not strongly dynamically coupled.

5.3. Dynamical stability

In order to validate the derived planetary architecture of the system, we performed a set of dynamical simulations to investigate the dynamical integrity of the orbits and possible eccentricity variations induced by the proximity of the planets to each other.

We performed one simulation including only planets b and c, and a set of different simulations on a range of possible values for the semi-major axes of planet d, given its uncertain location. We adopted the stellar parameters shown in Table 2 and the planetary properties provided by Table 3.5.

All simulations were carried out using the N-body integration capabilities of the Mercury Integrator Package (Chambers 1999) and adopting a methodology similar to the one described by Kane & Raymond (2014); Kane (2016). They were carried out on a hybrid symplectic/Bulirsch-Stoer integrator with a Jacobi coordinate system (Wisdom & Holman 1991; Wisdom 2006), with a time resolution of 0.2 days to ensure adequate sampling of the planetary orbits, on a time span of 10^7 years.

Results from the simulation of planets b and c alone show that they are in exceptionally stable orbits that remain approximately circular, inferring minimal dynamical interactions between them.

To study the effects of planet d, we assumed a value for the semi-major axes in the range from 0.19 to 0.26 AU, corresponding to the mean and -1σ values for the constrained period of 46 ± 4 days. We only considered the -1σ (and not the $+1\sigma$) scenario to put ourselves in the most dynamically extreme situation for which the planet orbits as close as possible (within our predictions) to the other two celestial bodies and its gravitational influence has a larger effect. The system remained stable through the semi-major axis range for planet d and for the full duration of the simulations. We further investigated the effects of high eccentricities for planet d and found that eccentricities as high as 0.4 still allow for fully stable orbits within the system even in the most compact configuration.

For example, Figure 10 shows the eccentricity evolution of the three planets for the extreme scenario where planet d is lo-

cated in a 0.19 AU orbit with a starting eccentricity of 0.4. Remarkably, the stability of the system enables a seamless transfer of angular momentum between the planets in long-term sustainable cycles. These results show that our understanding of the HD 22946 planetary system is dynamically sustainable, strengthening the self-consistency of our analysis.

5.4. Possible undetected planets

Here, we investigate the presence of potentially undetected planets orbiting in the gaps between the planets presented in this work following the same approach of Humphrey & Quintana (2020). Gladman (1993) defined a system of two planets as Hillstable (unstable) if Δ , i.e. the separation between the two, normalized by the mutual Hill-radius (R_H) of the pair, is above (below) an analytically defined critical level, Δ_{crit} . In the more general case of multi-planetary systems, Chambers et al. (1996); Pu & Wu (2015) used numerical simulations to retrieve the minimum value of Δ_{crit} above which the pair can be considered dynamically stable. Chambers et al. (1996) simulated systems with circular, coplanar orbits, and found that a $\Delta_{crit} = 10$ can be generally used to measure the stability of multi-planet systems within the described prescriptions. Pu & Wu (2015) found the same Δ_{crit} as Chambers et al. (1996) for systems on circular and coplanar orbits but extended their simulations by including systems eccentricities and inclinations drawn from a Rayleigh distribution. They find that planet pairs with $\Delta < \Delta_{crit} = 12.3$ are expected to experience close encounters, orbit crossing events and/or ejections due to strong gravitational interaction between the planets. Considering these two values for Δ_{crit} , we can check if the HD 22946 system could host, between each planet pair, an additional planet without jeopardizing the system stability. Following Humphrey & Quintana (2020), we computed the interval ($[a_{xmin}, a_{xmax}]$) of possible semi-major axes for a potential additional planet in a range of possible masses M_x for both the b/c and the c/d pair. If, for a given mass, $a_{xmin} \leq a_{xmax}$, the planet pair is considered unpacked, i.e. there is enough space to host an additional planet that would not destabilize the system. By solving equations (2) and (3) of Humphrey & Quintana (2020), one can compute (M_{xmax}, a_x) , the maximum mass that a potential undetected planet could have without destabilizing the system and its related semi-major axis. We computed a_{xmin} and a_{xmax} for each value of Δ_{crit} for both couples in the system. The




Fig. 11. Mass and semi-major axis of a possible undetected planet between TOI-411 b/c (upper panel) and TOI-411c/d (lower panel). Minimum semi-major axis (a_{xmin}) in dashed line and maximum one (a_{xmax}) in continuous line for $\Delta_{crit} = 10$ and 12.3 in red and green respectively. Given Δ_{crit} , the shaded region contains the possible combinations (M_x, a_x) for a stable system configuration hosting one more planet.

Pair b/c	M_{xmax}/M_{\oplus}	$a_x(AU)$
$\Delta_{crit} = 10$	18.6	0.0698
$\Delta_{crit} = 12.3$	6.7	0.0693
Pair c/d	M_{xmax}/M_{\oplus}	$a_x(AU)$
$\Delta_{crit} = 10$	127.4	0.156
$\Delta_{crit} = 12.3$	64.2	0.156

Table 5. Maximum mass M_{xmax} and related semi-major axis a_x of a potential additional planet orbiting between planets b/c or c/d. Two different values for Δ_{crit} are considered (see text).

results are shown in Figure 11. Both planet pairs are unpacked for each value of Δ_{crit} , though there is quite limited space for an additional planet between TOI-411 *b* and *c*. The TOI-411 *c* and *d*, on the other hand, might host an additional planet as large as 0.4 M_J based on the least stringent Δ_{crit} value of 10. Table 5 shows the computed values for M_{xmax} and related a_x under the two different Δ_{crit} conditions. We searched the light curve with the TLS pipeline in the intermediate range of periods and found no evidence for additional transiting candidates. However, there is the possibility that a potentially undetected planet is not observed transiting the star due to its orbital inclination and might be discovered via precise radial velocity techniques as in, e.g., Demangeon et al. (2021).

5.5. Potential for atmospheric characterization

The observed period and derived radius for TOI-411 b place it in the hot super-Earths range. Specifically, in the $1.5 - 2.0R_{\oplus}$ interval where a scarcity of exoplanets has been observed by Fulton et al. (2017). This is considered to be a transition region from rocky worlds, with a high-molecular-weight atmosphere, to lowdensity worlds, dominated by a H/He gaseous envelope. The former may have accreted a primary atmosphere in the early stages of formation that might have been dispersed due to escape processes, such as stellar radiation photo-evaporation (Fulton & Petigura 2018) or core-powered mass loss (Ginzburg et al. 2018). The only super-Earth with a detected (and largely discussed) atmosphere is 55 Cnc e, which probably lost its primary H/He envelope (Zhang et al. 2021). Even if TOI-411 b might have lost its primary atmosphere during its early evolution, a large reserve of hydrogen could still be trapped in its mantle (Kite et al. 2019). This could lead to the out-gassing of a secondary atmosphere, as suggested in the case of the Earth-like planet GJ 1132 b (Swain et al. 2021). Precise radial velocity measurements and improved mass estimates will be critical to determining the mean density of the planet and constraining the planet's composition. TOI-411 c and the single transit event candidate appear to be, respectively, a hot and a warm sub-Neptune, hence they could potentially hold a thick H/He envelope. Future observations with new generation telescopes of these < 4 R_{\oplus} exoplanets, will be crucial to identify their atmospheric composition and narrow down the possibilities regarding their formation mechanisms. For this purpose, following the methods of Kempton et al. (2018), we estimated the potential for atmospheric follow-up of these worlds with the James Webb Space Telescope (JWST). We computed the Transmission Spectroscopy Metric (TSM), a quantity proportional to the expected SNR of the transmission spectrum of a cloud-free atmosphere, over a 10-hour observation in the NIRISS bandpass. The TSM values for planets b and c and candidate d are, respectively, 65 ± 10 , 89 ± 16 and 67 ± 14 . We note that the TSM value is intended as a rough indicator for transmission spectroscopy follow-ups that can and should be refined with precise mass measurement of the planets. We put our planets into context by comparing their TSM with the same quantity for similar worlds. Specifically, Fig.12 shows the TSM values for every confirmed planet listed on the NASA Exoplanet Archive¹³ orbiting F/G stars, with $1.5R_{\oplus} < R_p < 4.0R_{\oplus}$ and $P_{orb} < 50d$. The lower limit is chosen to discard the terrestrial planets for which the Kempton et al. (2018) TSM threshold is set to 10 whilst the upper one keeps the sample within the sub-Neptune radius range. The reported equilibrium temperature (T_{eq}) has been computed for each planet by assuming null albedo and full day-night heat redistribution. Within this sample of 1140 planets, only 1.2% of them have a TSM value above the threshold of 90 that Kempton et al. (2018) indicates to qualify the planet as suitable for atmospheric detection with JWST, and TOI-411 c can potentially overcome this threshold.

6. Conclusions

We present the discovery of a multi-planetary system around the bright ($V \sim 8.3$ mag), nearby (~ 63 pc) Sun-like star HD 22946, indicated also as TOI-411, from *TESS* data. The host star was observed by *TESS* in Sectors 3,4, 30 and 31 and two transiting planets with periods of 4.04 and 9.57 days were identified in the light curve, with an additional single event that we associate to a

¹³ https://exoplanetarchive.ipac.caltech.edu/index.html




Fig. 12. Transmission Spectroscopy Metric (TSM) values for known exoplanets orbiting F/G stars in less than 50 days. We only show exoplanets for which TSM> 50 for clarity as ~ 98% of worlds in the sample lays below this value. While TOI-411 *b* and *c* fall in the 2% of this kind of planet with TSM> 50, only TOI-411 *c* passes the suggested threshold of 90, joining an elite group of 15 super-Earths and sub-Neptunes orbiting Sun-like stars amenable for spectroscopic follow-ups with JWST.

third transiting planet with a period of 46 ± 4 days. By modeling the transits, we determine the planets radii: $1.74 \pm 0.10 R_{\oplus}$ for planet *b*, $2.74 \pm 0.14 R_{\oplus}$ for planet *c* and $3.23 \pm 0.19 R_{\oplus}$ for planet *d*.

While the nature of the single event cannot be verified by using exclusively *TESS* data and follow-ups are not available yet, the innermost exoplanets have been vetted under several different aspects: the pipelines DAVE, vespa and triceratops have been used to exclude false positive scenarios or determine that the false positive probability is negligible, whilst TFOP photometric data and high-resolution imaging have been used to rule out faint bound companions and foreground/background or nearby eclipsing binaries that could contaminate the *TESS* light curve.

Therefore, we validated the discovery of planets b (or TOI-411.02) and c (or TOI-411.01), while we plan to collect additional data (radial-velocity curve and photometric follow-up observations) in order to further confirm the nature of candidate planet d (or TOI-411.03)

Multi-planetary systems around Sun-like stars, such as the one reported here, are ideal laboratories to test our understanding of how planets form end evolve. TOI-411 *b*, with an inferred mass of ~ 3.5 M_{\oplus} , is a hot super-Earth (Fulton et al. 2017; Fressin et al. 2013), orbiting its star with a period of \approx 4 days at the equilibrium temperature of 1378±36 K. Because of its size, it is one of the relatively few planets found within the Fulton gap, i.e., a range of planetary radii between 1.5 and $2R_{\oplus}$ that is relatively less populated (Fulton et al. 2017) with respect to smaller or larger radii. This bimodality in the planets' distribution is thought to be the effect of photoevaporation of volatile gases or core-powered mass loss that strips low-mass planets of

their atmosphere, leaving behind bare, rocky planets, while gas giants with radii larger than $2R_{\oplus}$ remain unaffected.

Indeed, TOI-411 c and TOI-411 d have larger radii, with a radius ratio of 1.6 and 1.8 respectively. By adopting the approach described in Humphrey & Quintana (2020), we find that between planet b and c it would be possible to include at least another planet, without compromising the dynamical stability of the system. In the scenario described above, this undetected planet would have an intermediate period and a smaller radius ratio. However, we did not find evidence for this intermediate planet from the *TESS* light curve.

With an equilibrium temperature of about 1000 K, planet *c* is a hot sub-Neptunian, and holds particular interest because it *can* potentially host a tick atmosphere and has a TSM compatible with atmospheric detection from JWST. In fact, even if the host star is bright, its luminosity (~ 7.3 mag in the *J*-band) is still below the saturation limit for the NIRISS single-object spectroscopy mode. An independent constraint on the mass value for this planet from radial-velocity measurement is critical to confirm the suitability of this target for atmospheric detection. On the other side, by being a cold (T_{eq} =622 K) sub-Neptunian, planet *d* would provide the opportunity of making comparative atmospheric characterization via transmission spectroscopy, if observable. In its case, the TSM estimate should be recomputed when the validation is confirmed and the period is known with better precision.

In summary, the HD 22946 multiplanetary system offers many interesting insights into planetary formation, by hosting a star that is bright and similar to our Sun, three planets with a common origin but a completely different aspect: one small planet stripped of its atmosphere (or with a secondary, accreted atmosphere), two larger planets at significantly different temperature, and probably yet-to-be-discovered intermediate ones.

Therefore, this system provides an exciting opportunity for a synergy of space- and ground-based facilities such as the CHaracterising ExOplanets Satellite (CHEOPS Broeg et al. 2013), HARPS and ESPRESSO to determine the bulk density of the planets, and, possibly, JWST to investigate the presence of the atmosphere. This will help us to gain a better understanding of the formation pathways that can lead to a similar system.

Acknowledgements. We thank the referee for her/his comments that helped to improve and clarify the presentation of our results. This paper includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA's Science Mission Directorate. We acknowledge the use of public TESS Alert data from pipelines at the TESS Science Office and at the TESS Science Processing Operations Center. This research has made use of the Exoplanet Follow-up Observation Program website, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center for the production of the SPOC data products. L.I. acknowledges the "PON Ricerca e Innovazione: Attraction and International Mobility (AIM)" program for support. This work has made use of data from the European Space Agency (ESA) mission Gaia (https://www.cosmos.esa.int/gaia), processed by the Gaia Data Processing and Analysis Consortium (DPAC, https://www.cosmos.esa.int/web/gaia/dpac/consortium). Observations in the paper made use of the High-Resolution Imaging instrument Zorro at Gemini-South. The data underlying this article is publicly available at the following archives: MAST¹⁴ and ExoFOP-TESS¹⁵.

This work is based on data collected at the following *facilities*: ASAS-SN, Exoplanet Archive, *Gaia*, Gemini:South (Zorro), MAST, LCOGT, SOAR (HRcam), TESS.

This work makes use of the following *software*: astropy, lightkurve (Lightkurve Collaboration et al. 2018), transitleastsquares (Hippke & Heller 2019), DAVE

Article number, page 15 of 21

¹⁴ https://archive.stsci.edu/missions-and-data/tess

¹⁵ https://exofop.ipac.caltech.edu/tess/

(Kostov et al. 2019), exoplanet [(Foreman-Mackey et al. 2017),(Foreman-Mackey 2018)], vespa (Morton 2015), triceratops (Giacalone & Dressing 2020), tpfplotter (Aller et al. 2020).

References

- 1997, ESA Special Publication, Vol. 1200, The HIPPARCOS and TYCHO catalogues. Astrometric and photometric star catalogues derived from the ESA HIPPARCOS Space Astrometry Mission
- Aller, A., Lillo-Box, J., Jones, D., Miranda, L. F., & Barceló Forteza, S. 2020, A&A, 635, A128
- Bitsch, B. 2019, A&A, 630, A51 Broeg, C., Fortier, A., Ehrenreich, D., et al. 2013, in European Physical Journal Web of Conferences, Vol. 47, European Physical Journal Web of Conferences, 03005
- Brown, T. M., Baliber, N., Bianco, F. B., et al. 2013a, PASP, 125, 1031
- Brown, T. M., Baliber, N., Bianco, F. B., et al. 2013b, PASP, 125, 1031
- Buchhave, L. A., Latham, D. W., Johansen, A., et al. 2012, Nature, 486, 375
- Cacciapuoti, L., Kostov, V. B., Kuchner, M., et al. 2022, MNRAS, 513, 102
- Chambers, J., Wetherill, G., & Boss, A. 1996, Icarus, 119, 261
- Chambers, J. E. 1999, MNRAS, 304, 793
- Chen, J. & Kipping, D. 2017, ApJ, 834, 17
- Ciardi, D. R., Beichman, C. A., Horch, E. P., & Howell, S. B. 2015, ApJ, 805, 16 Collins, K. 2019, in American Astronomical Society Meeting Abstracts, Vol.
- 233, American Astronomical Society Meeting Abstracts #233, 140.05 Collins, K. A., Kielkopf, J. F., Stassun, K. G., & Hessman, F. V. 2017, AJ, 153,
- 77
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, VizieR Online Data Catalog, II/246
- Demangeon, O. D. S., Zapatero Osorio, M. R., Alibert, Y., et al. 2021, A&A, 653, A41
- Donati, J.-F., Semel, M., Carter, B. D., Rees, D. E., & Collier Cameron, A. 1997, MNRAS, 291, 658
- Eisner, N. L., Barragán, O., Lintott, C., et al. 2021, MNRAS, 501, 4669
- Fabrycky, D. C., Lissauer, J. J., Ragozzine, D., et al. 2014, ApJ, 790, 146
- Faigler, S. & Mazeh, T. 2011, MNRAS, 415, 3921
- Foreman-Mackey, D. 2018, Research Notes of the American Astronomical Society, 2, 31
- Foreman-Mackey, D., Agol, E., Ambikasaran, S., & Angus, R. 2017, AJ, 154, 220
- Foreman-Mackey, D., Luger, R., Agol, E., et al. 2021, The Journal of Open Source Software, 6, 3285
- Fressin, F., Torres, G., Charbonneau, D., et al. 2013, ApJ, 766, 81
- Fulton, B. J. & Petigura, E. A. 2018, AJ, 156, 264
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, AJ, 154, 109
- Furlan, E., Ciardi, D. R., Everett, M. E., et al. 2017, AJ, 153, 71
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2021, A&A, 650, C3 Giacalone, S. & Dressing, C. D. 2020, triceratops: Candidate exoplanet rating
- tool
- Giacalone, S., Dressing, C. D., Jensen, E. L. N., et al. 2021, AJ, 161, 24
- Gillon, M., Triaud, A. H. M. J., Demory, B.-O., et al. 2017, Nature, 542, 456
- Ginzburg, S., Schlichting, H. E., & Sari, R. 2018, MNRAS, 476, 759
- Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, A&AS, 141, 371
- Gladman, B. 1993, Icarus, 106, 247
- Guerrero, N. & TESS Science Office. 2021, in American Astronomical Society Meeting Abstracts, Vol. 53, American Astronomical Society Meeting Abstracts, 117.01
- Hippke, M. & Heller, R. 2019, A&A, 623, A39
- Hoffman, M. D. & Gelman, A. 2014, Journal of Machine Learning Research, 15, 1593
- Howell, S. B., Everett, M. E., Horch, E. P., et al. 2016, ApJ, 829, L2
- Howell, S. B., Everett, M. E., Sherry, W., Horch, E., & Ciardi, D. R. 2011, AJ, 142, 19
- Huang, C. X., Vanderburg, A., Pál, A., et al. 2020, Research Notes of the American Astronomical Society, 4, 204
- Humphrey, A. L. T. & Quintana, E. V. 2020, arXiv preprint arXiv:2011.03053
- Jayasinghe, T., Kochanek, C. S., Stanek, K. Z., et al. 2021, MNRAS, 503, 200 Jenkins, J. M. 2002, ApJ, 575, 493
- Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9913, Software and Cyberinfrastructure for Astronomy IV, ed. G. Chiozzi & J. C. Guzman, 99133E
- Jensen, E. 2013, Tapir: A web interface for transit/eclipse observability, Astrophysics Source Code Library
- Kane, S. R. 2016, ApJ, 830, 105

Kane, S. R. & Raymond, S. N. 2014, ApJ, 784, 104

Article number, page 16 of 21

Kempton, E. M. R., Bean, J. L., Louie, D. R., et al. 2018, PASP, 130, 114401

- Kervella, P., Arenou, F., Mignard, F., & Thévenin, F. 2019, A&A, 623, A72
- Kervella, P., Arenou, F., & Thévenin, F. 2022, A&A, 657, A7
- Kervella, P., Thévenin, F., Di Folco, E., & Ségransan, D. 2004, A&A, 426, 297 Kipping, D. M. 2013a, MNRAS, 435, 2152
- Kipping, D. M. 2013b, MNRAS, 435, 2152
- Kite, E. S., Fegley Jr, B., Schaefer, L., & Ford, E. B. 2019, The Astrophysical Journal Letters, 887, L33
- Kostov, V. B., Mullally, S. E., Quintana, E. V., et al. 2019, AJ, 157, 124
- Kovács, G., Zucker, S., & Mazeh, T. 2002, A&A, 391, 369
- Latham, D. W., Rowe, J. F., Quinn, S. N., et al. 2011, ApJ, 732, L24
- Li, J., Tenenbaum, P., Twicken, J. D., et al. 2019, Publications of the Astronomical Society of the Pacific, 131, 024506
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, Lightkurve: Kepler and TESS time series analysis in Python, Astrophysics Source Code Library
- Lissauer, J. J., Marcy, G. W., Bryson, S. T., et al. 2014, ApJ, 784, 44 Lissauer, J. J., Marcy, G. W., Rowe, J. F., et al. 2012, ApJ, 750, 112
- Lissauer, J. J., Ragozzine, D., Fabrycky, D. C., et al. 2011, ApJS, 197, 8
- Lomb, N. R. 1976, Ap&SS, 39, 447
- Loyd, R. O. P., Shkolnik, E. L., Schneider, A. C., et al. 2020, ApJ, 890, 23
- Luger, R., Agol, E., Foreman-Mackey, D., et al. 2019, AJ, 157, 64
- Luhn, J. K., Wright, J. T., Howard, A. W., & Isaacson, H. 2020, AJ, 159, 235
- Mamajek, E. E. & Hillenbrand, L. A. 2008, ApJ, 687, 1264
- McCully, C., Volgenau, N. H., Harbeck, D.-R., et al. 2018a, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 10707, Software and Cyberinfrastructure for Astronomy V, ed. J. C. Guzman & J. Ibsen, 107070K
- McCully, C., Volgenau, N. H., Harbeck, D.-R., et al. 2018b, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 10707, Proc. SPIE, 107070K
- Mermilliod, J. C. 2006, VizieR Online Data Catalog, II/168
- Miret-Roig, N., Bouy, H., Raymond, S. N., et al. 2021, arXiv e-prints, arXiv:2112.11999
- Morris, R. L., Twicken, J. D., Smith, J. C., et al. 2020, Kepler Data Processing Handbook: Photometric Analysis, Kepler Science Document KSCI-19081-003
- Morris, S. L. & Naftilan, S. A. 1993, ApJ, 419, 344
- Morton, T. D. 2015, VESPA: False positive probabilities calculator
- Murgas, F., Jenkins, J. S., Rojo, P., Jones, H. R. A., & Pinfield, D. J. 2013, A&A, 552, A27
- Naef, D., Latham, D. W., Mayor, M., et al. 2001, A&A, 375, L27
- Otegi, J. F., Bouchy, F., & Helled, R. 2020, A&A, 634, A43 Owen, J. E. & Wu, Y. 2017, ApJ, 847, 29

- Paredes, L. A., Henry, T. J., Quinn, S. N., et al. 2021, AJ, 162, 176 Pepe, F., Mayor, M., Rupprecht, G., et al. 2002, The Messenger, 110, 9
- Pepe, F., Molaro, P., Cristiani, S., et al. 2014, Astronomische Nachrichten, 335, 8
- Percy, J. R., Au-Yong, K., Gilmour-Taylor, G., et al. 2002, in Astronomical Society of the Pacific Conference Series, Vol. 256, Observational Aspects of Pulsating B- and A Stars, ed. C. Sterken & D. W. Kurtz, 99
- Petigura, E. A., Howard, A. W., Marcy, G. W., et al. 2017, AJ, 154, 107
- Pojmanski, G. 1997, Acta Astron., 47, 467
- Pribulla, T., Sebastian, D., Ammler-von Eiff, M., et al. 2014, MNRAS, 443, 2815
- Pu, B. & Wu, Y. 2015, The Astrophysical Journal, 807, 44
- Ragozzine, D. & Holman, M. J. 2010, arXiv e-prints, arXiv:1006.3727
- Richards, J. W., Starr, D. L., Miller, A. A., et al. 2012, ApJS, 203, 32
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003
- Rodriguez, J. E., Quinn, S. N., Zhou, G., et al. 2021, AJ, 161, 194
- Rowe, J. 2016, Kepler: Kepler Transit Model Codebase Release.
- Rowe, J. F., Bryson, S. T., Marcy, G. W., et al. 2014, ApJ, 784, 45
- Rowe, J. F. & Thompson, S. E. 2015, arXiv e-prints, arXiv:1504.00707
- Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. 2016, PeerJ Computer Science, 2, e55

Siverd, R. J., Brown, T. M., Barnes, S., et al. 2018, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 10702, Ground-

Siverd, R. J., Brown, T. M., Hygelund, J., et al. 2016, in Society of Photo-Optical

based and Airborne Instrumentation for Astronomy VII, ed. C. J. Evans,

Instrumentation Engineers (SPIE) Conference Series, Vol. 9908, Ground-

based and Airborne Instrumentation for Astronomy VI, ed. C. J. Evans,

Scargle, J. D. 1982, ApJ, 263, 835

Shporer, A. 2017, PASP, 129, 072001

L. Simard, & H. Takami, 107026C

L. Simard, & H. Takami, 99086X

Space Sciences, 8, 138

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525 Scott, N. J., Howell, S. B., Gnilka, C. L., et al. 2021, Frontiers in Astronomy and

Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, ApJ, 788, 48

Seager, S. & Mallén-Ornelas, G. 2003, ApJ, 585, 1038

Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163 ¹ Department of Physics "Ettore Pancini", University of Naples, Federico II 2 European Southern Observatory, Karl-Schwarzschild-Strasse 2 D-Stassun, K. G. & Torres, G. 2016, AJ, 152, 180 85748 Garching bei Munchen, Germany Science and Technology Department, Parthenope University of Naples, CDN IC4, 80143, Naples, Italy Stassun, K. G. & Torres, G. 2018, ApJ, 862, 61 INAF-Osservatorio Astronomico di Capodimonte, Salita Moraliello, 16, 80131, Naples, Italy 5 INFN, Sezione di Napoli, Italy ⁶ NASA Goddard Space Flight Center, Exoplanets and Stellar Astro-Stassun, K. G. & Torres, G. 2021, ApJ, 907, L33 physics Laboratory (Code 667), Greenbelt, MD 20771, USA Vanderbilt University, Department of Physics & Astronomy, 6301 Stevenson Center Lane, Nashville, TN 37235, USA Fisk University, Department of Physics, 1000 18th Avenue N., Stumpe, M. C., Smith, J. C., Catanzarite, J. H., et al. 2014, PASP, 126, 100 Nashville, TN 37208, USA Department of Astronomy, University of Maryland, College Park, MD 20742, USA Swain, M. R., Estrela, R., Roudier, G. M., et al. 2021, The Astronomical Journal, 10 Department of Astronomy, University of California Berkeley, 161, 213 Berkeley, CA 94720-3411, USA 11 Department of Earth and Planetary Sciences, University of California, Riverside, CA 92521, USA 12 SETI Institute, 189 Bernardo Ave, Suite 200, Mountain View, CA Tokovinin, A. 2018a, PASP, 130, 035002 94043. USA 13 Department of Physics and Astronomy, Bishops University 2600 College St, Sherbrooke, QC J1M 1Z7, Canada Tokovinin, A. 2018b, PASP, 130, 035002 Tsinghua International School, Beijing 100084, China 15 George Mason University, 4400 University Drive, Fairfax, VA, 22030 USA 16 Center for Astrophysics |Harvard & Smithsonian, 60 Garden Street, Torres, G., Andersen, J., & Giménez, A. 2010, A&A Rev., 18, 67 Cambridge, MA 02138, USA 17 Perth Exoplanet Survey Telescope, Perth, Western Australia 18 Departamento de Matematica y Fisica Aplicadas, Facultad de Ingegneria, Universidad Catolica de la Santisima Concepcion, Alonso de Twicken, J. D., Catanzarite, J. H., Clarke, B. D., et al. 2018, PASP, 130, 064502 Rivera 2850, Concepcion, Chile 19 Caltech/IPAC-NASA Exoplanet Science Institute, 770 S. Wilson Avenue, Pasadena, CA 91106, USA Twicken, J. D., Chandrasekaran, H., Jenkins, J. M., et al. 2010, in Society 20 NASA Exoplanet Science Institute, Caltech/IPAC, Mail Code 100of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 22, 1200 E. California Blvd., Pasadena, CA 91125, USA 7740, Software and Cyberinfrastructure for Astronomy, ed. N. M. Radziwill NASA Ames Research Center, Moffett Field, CA 94035, USA & A. Bridger, 77401U ²² Department of Physics, Engineering and Astronomy, Stephen F. Austin State University, 1936 North St, Nacogdoches, TX 75962, USA 23 Cerro Tololo Inter-American Observatory, Casilla 603, La Serena, VanderPlas, J. T. 2018, ApJS, 236, 16 Chile 24 Department of Physics and Astronomy, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3255, USA Winn, J. N., Matthews, J. M., Dawson, R. I., et al. 2011, ApJ, 737, L18 Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA3902139, USA ²⁶ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Wisdom, J. 2006, AJ, 131, 2294 Cambridge, MA 02138, USA Institute of Space Astrophysics and Planetology INAF-IAPS, Via Fosso del Cavaliere 100, I-00133, Rome, Italy Wisdom, J. & Holman, M. 1991, AJ, 102, 1528 28 Western Connecticut State University, 181 White Street, Danbury CT, 06810. 29 Department of Astronomy and Carl Sagan Institute, Cornell University, 122 Sciences Drive, Ithaca, NY 14853, USA Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, AJ, 140, 1868 30 Proto-Logic LLC, 1718 Euclid Street NW, Washington, DC 20009, USA

31

32

33

34

35

Zhang, M., Knutson, H. A., Wang, L., et al. 2021, The Astronomical Journal, 161, 181

Ziegler, C., Tokovinin, A., Briceño, C., et al. 2020, AJ, 159, 19

Ziegler, C., Tokovinin, A., Latiolais, M., et al. 2021, AJ, 162, 192

³⁷ University of Southern Queensland, <u>Contere four Astrophysics</u>, West Street, Toowoomba, QLD 4350 Australia

Department of Earth, Atmospheric and Planetary Sciences, Mas-

Kavli Institute for Astrophysics and Space Research, Massachusetts

Department of Astrophysical Sciences, Princeton University,

Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA Department of Aeronautics and Astronautics, Massachusetts Insti-

sachusetts Institute of Technology, Cambridge, MA 02139, USA

Institute of Technology, Cambridge, MA 02139, USA

SETI Institute, Mountain View, CA 94043, USA

tute of Technology, Cambridge, MA 02139, US

Princeton, NJ 08544, USA

- ³⁸ Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ³⁹ Curtin Institute of Radio Astronomy, Curtin University, Bentley, Western Australia 6102
 - D (CDI ' LA CAT M'

Appendix A: DAVE results

In this Section we show examples of DAVE results. Figures A.1 and A.2 show the photocenter difference images for the planet candidates as computed in different sectors. The black star indicates the TIC position, the red circle is the average of measured photocenters for each transit. The white dashed line indicates the TESS target pixel aperture used to extract the light curve. Figures A.3 and A.4 show TESS transit data (first panel), the binned data (second panel), diagnostic plots to check for odd-even effects (third panel) secondary, tertiary eclipses and positive bumps, as might be produced by false positives (bottom panel). DAVE analysis reports no significant false positive indicators for TOI 411 b and c.




Fig. A.1. DAVE photocenter out-of-transit, in-transit, difference and difference SNR images for planet b based on TESS sector 3 (upper panel) and 30 (upper panel) data. The axes numbers are expressed in TESS pixels.




Fig. A.2. DAVE photocenter out-of-transit, in-transit, difference and difference SNR images for planet c based on sector 30 (upper panel) and 31 (lower panel) data. The axes numbers are expressed in TESS pixels.




Fig. A.3. DAVE Modelshift for planet b based on TESS data of Sector 30. No significant false positive indicator is detected.

Cacciapuoti et al.: A multiplanetary system around HD 22946




Fig. A.4. DAVE Modelshift for planet c based on TESS data of Sector 31. No significant false positive indicator is detected.